



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

XXI International Grassland Congress / VIII  
International Rangeland Congress

---

## Effects of Different Salts and Concentrations on Germination and Growth of *Leymus chinensis*

L. H. Huang

*Chinese Academy of Sciences, China*

Z. W. Liang

*Chinese Academy of Sciences, China*

H. Y. Ma

*Chinese Academy of Sciences, China*

M. Liu

*Chinese Academy of Sciences, China*

C. Yan

*Chinese Academy of Sciences, China*

*See next page for additional authors*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/14-1/3>

The XXI International Grassland Congress / VIII International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

---

### **Presenter Information**

L. H. Huang, Z. W. Liang, H. Y. Ma, M. Liu, C. Yan, and Z. C. Wang

## Effects of different salts and concentrations on germination and growth of *Leymus chinensis*

L.H. Huang, Z.W. Liang\*, H.Y. Ma, M. Liu, C. Yan, Z.C. Wang

Northeast Institute of Geography and Agricultural Ecology, CAS, Changchun, Jilin 130012, China; Da'an Sodic Land Experiment Station of China, Da'an Jilin 131317, China; \*E-mail: liangzw@neigae.ac.cn

**Key words:** sodium salt, potassium salt, stress, germination, growth, *Leymus chinensis*

**Introduction** *Leymus chinensis* is one of good-quality and saline-alkali tolerant grasses in the northern area of China with economic and ecological significance. The high concentration of salt ions is the main factor inhibiting *L. chinensis* seed germination in the sodic soil in Songnen plain, while the level of inhibition depends upon the concentrations and kinds of salt. This study is to investigate the effect of sodium salts ( $\text{NaCl}$ ,  $\text{Na}_2\text{SO}_4$ ,  $\text{NaHCO}_3$ ,  $\text{Na}_2\text{CO}_3$ ) and potassium salts ( $\text{KCl}$ ,  $\text{K}_2\text{SO}_4$ ,  $\text{KHCO}_3$ ,  $\text{K}_2\text{CO}_3$ ) on germination of *L. chinensis*.

**Materials and methods** In this experiment, 0, 5, 10, 20, 50, and 100 mM  $\text{NaCl}$ ,  $\text{Na}_2\text{SO}_4$ ,  $\text{NaHCO}_3$ ,  $\text{Na}_2\text{CO}_3$ ,  $\text{KCl}$ ,  $\text{K}_2\text{SO}_4$ ,  $\text{KHCO}_3$ ,  $\text{K}_2\text{CO}_3$  were prepared and pH and electrical conductivity (EC) of solutions were determined. Seeds were placed on the plastic web floating upon 200 ml solution in a beaker and the solution was added every 2~3 d. The germination temperature was 16/28°C with 12 h dark/12 h light photoperiod and the germination percentage of seeds, length of root and shoot were calculated and measured after 21 days.

**Results** In the eight salt solutions, pH of  $\text{Na}_2\text{CO}_3$  and  $\text{K}_2\text{CO}_3$  was the highest (10.5~11.5), followed by  $\text{NaHCO}_3$  and  $\text{KHCO}_3$  (8.0~8.5). pH of the other 4 salts was the lowest at 6.5~7.0. The EC showed upward trend with the increasing salt concentrations. Germination percentage and length of root and shoot of *L. chinensis* decreased as the salt concentrations increased (Table 1). The inhibition of carbonate and bicarbonate on seed germination was the greatest and no germination occurred in  $\text{Na}_2\text{CO}_3$  or  $\text{K}_2\text{CO}_3$  at 100 mM.

**Table 1** Effect of different concentrations of eight salts on seed germination and growth of *Leymus chinensis*.

Salt kinds	Salt concentrations (mM)														
	5	10	20	50	100	5	10	20	50	100	5	10	20	50	100
	Germination percentage					Root length					Shoot length				
<i>NaCl</i>	↑	↑	↑	▽	▼	▽	▼	▼	▼	▼	↑	↑	↑	▼	▼
<i>KCl</i>	↑	↑	↑	▽	▼	▽	▼	▼	▼	▼	↑	↑	↑	▽	▼
<i>Na<sub>2</sub>SO<sub>4</sub></i>	↑	▽	▽	▼	▼	▼	▼	▼	▼	▼	↑	▽	▼	▼	▼
<i>K<sub>2</sub>SO<sub>4</sub></i>	↑	↑	▽	▽	▼	▼	▼	▼	▼	◎	↑	↑	▼	▼	▼
<i>NaHCO<sub>3</sub></i>	▽	▽	▼	▼	▼	↑	▼	▼	▼	◎	↑	↑	↑	▼	▼
<i>KHCO<sub>3</sub></i>	▽	▽	▽	▼	▼	▼	▼	▼	◎	◎	↑	↑	▼	▼	▼
<i>Na<sub>2</sub>CO<sub>3</sub></i>	↑	▽	▼	▼	◎	▼	▼	◎	◎	◎	↑	▽	▼	▼	◎
<i>K<sub>2</sub>CO<sub>3</sub></i>	↑	↑	▽	▼	◎	▼	▼	◎	◎	◎	↑	▼	▼	▼	◎

\* ↑ Stimulate germination slightly; ▽ Inhibit germination slightly; ▼ Inhibit germination markedly; ◎ Inhibit germination absolutely.

**Conclusion** These results indicated that pH of  $\text{Na}_2\text{CO}_3$  and  $\text{K}_2\text{CO}_3$  was high, which inhibit the germination and growth of root and shoot of *L. chinensis* even at lower concentrations.

**Acknowledgement** This project is supported by the 973 program (2007CB106803), National Key Project for the Eleventh Five Year Plan (2006BAC01A08), and the Foundation of the Knowledge Innovation Project of Chinese Academy of Sciences (No. KZCX3-SW-NA3-05).